

AD \_\_\_\_\_


GRANT NO: DAMD 17-93-J-3004

TITLE: Custom Tailoring therapy for breast cancer using erbB-2

PRINCIPAL INVESTIGATOR: Soonmyoung Paik, M.D.

CONTRACTING ORGANIZATION:  
Georgetown University  
37th & O Street, NW  
Washington, DC 20057

REPORT DATE: June 5, 1995



TYPE OF REPORT: Annual Report

PREPARED FOR: U.S. Army Medical Research and Materiel  
Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;  
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

DTIC QUALITY INSPECTED 5

19950823 024

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 5 June 95	3. REPORT TYPE AND DATES COVERED annual report (15 Dec 93-14 Dec 94)	
4. TITLE AND SUBTITLE Custom Tailoring Therapy for Breast Cancer using erbB-2			5. FUNDING NUMBERS DAMD 17-93-J-3004	
6. AUTHOR(S) Soonmyoung Paik, M.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Georgetown University Washington, DC 20057			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Since erbB-2 failed to be a therapeutic response varibale for tamoxifen response, we have used gene trap U3lacZ to clone estrogen regulated molecules in breast cancer cells. Bcl-2 gene was identified as an estrogen inducible gene. A novel estrogen repressed gene was cloned. Their potential role as therapeutic response variable is being examined. NSABP B-11 was examined for erbB-2 expression. Statistical analysis of the B-11 is pending.				
14. SUBJECT TERMS erbB-2, breast cancer, tamoxifen, gene trap, bcl-2, transcriptional repression			15. NUMBER OF PAGES 6	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

## FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the US Army.

X Where copyrighted material is quoted, permission has been obtained to use such material.

X Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

X Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

X In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

X For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

X In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

X In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

X In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Accession For	
NIH GRAAI	<input checked="" type="checkbox"/>
NIH TAB	<input type="checkbox"/>
Unpublished	<input type="checkbox"/>
Publication	
Distribution/	
Availability Codes	
Mail &/or	Special
A-1	

John Pa 6.30-95  
PI - Signature Date

## TABLE OF CONTENTS

Annual Report  
June 5, 1995  
Soonmyoung Paik, M.D.

Front Cover	Page 1
SF 298	Page 2
Foreword	Page 3
Table of Contents	Page 4
Summary	Page 5
Introduction	Page 5
Body	Page 6
Conclusion	Page 6

## Annual Report

Title: Custom tailoring of chemotherapy with erbB-2

### Summary:

Our initial proposal was to examine erbB-2 as a therapeutic response variable that predicts the response to specific chemo or hormonal therapy in the treatment of breast cancer. Initial phase of the project involved the examination of 937 cases from the protocol B-14 of National Surgical Adjuvant Breast Project (NSABP). In this study, the value of erbB-2 as well as other markers such as S-phase fraction in predicting response to tamoxifen was examined. Since as stated in the last annual report erbB-2 failed to predict response to tamoxifen, I have attempted to identify other markers that would have predictive value. Instead of examining other known markers, I elected to identify and clone new markers using retroviral gene trap. Using this approach, I have identified bcl-2 as an estrogen regulated molecule in breast cancer cells. In addition, a novel gene has been identified that is repressed upon estrogen treatment. Value of these new markers in the prediction of tamoxifen response will be examined in the future. Meanwhile, as a continuation of the originally proposed project on erbB-2, we have screened cases from NSABP B-11, in which patients were randomized to receive adriamycin vs. non-adriamycin adjuvant regimen. Unfortunately, due to the problems at the headquarters of NSABP, we have not yet completed the statistical analysis of these cases.

### Introduction:

Treatment of breast cancer has been revolutionized through three important stages. First was the development of surgical techniques to remove breast cancer tissue which improved the survival of patients dramatically. Second stage was the use of systemic adjuvant therapy (tamoxifen for estrogen receptor positive tumors and chemotherapy for estrogen receptor negative tumors). The final revolution was the early detection through screening program. Even after these improvements in the diagnosis and management, we still did not obtain 100% cure of breast cancer. Thus, about 30% of early invasive breast cancer will still recur within 10 years after surgery and radiation. Additional systemic therapy will reduce this recurrence rate to about 20%. This modest benefit brings significant dilemma for clinical practice. Thus only small subset of patients actually benefit from systemic therapy. Focus of our lab has been to identify and test markers that predict response to systemic therapy. Our early attention to erbB-2 as a potential marker for tamoxifen response stems from in-vitro studies that demonstrated negative interaction between estrogen receptor signalling pathway and erbB-2 pathway. However, using samples from NSABP B-14, we failed to see any impact of erbB-2 in tamoxifen response. Thus we have changed our attention to other potential markers that will be useful as a predictor of tamoxifen response.

One of the important development in the recent year or two has been the identification apoptosis as a key process in cancer development and in chemotherapeutic action mechanism. Thus molecules involved in apoptotic pathway are potential candidates for therapeutic response variables. Estrogen is a key survival factor for breast epithelial cells. We have postulated that estrogen should inhibit molecules that are involved in apoptosis in order to maintain cell survival.

Thus we elected to clone genes that are regulated by estrogen.

### **Body:**

Breast cancer cells and normal ductal epithelial cells which express the estrogen receptor (ER) undergo apoptosis upon estrogen withdrawal. This suggests the importance of estrogen in preventing apoptosis of estrogen receptor positive mammary ductal epithelial cells. For estrogen to suppress apoptosis it has to either repress molecular inducers of apoptosis and/or induce the expression of suppressors of apoptosis.

We have used the retroviral promoter trap U3lacZ to identify estrogen regulated genes from breast cancer cells so such genes could have an important role in the regulation of the apoptotic process in breast cancer cells. Clones in which the U3lacZ virus had integrated into genes which were either suppressed or induced by estrogen were identified. Beta-galactosidase reporter activity in 20 of 2000 clones mutated by U3lacZ were inducible by estrogen while 3 in 2000 were suppressed. Cloning and sequencing of 5'-flanking genomic DNA from one of the estrogen inducible clones showed 100% identity with 5'-untranslated region of published bcl-2 cDNA sequence. LacZ reporter activity in this clone showed the classical regulation pattern of an estrogen regulated gene. Bcl-2 is believed to play a key role in the apoptotic process in hormone dependent organs including mammary gland. The induction of bcl-2 by estrogen in the breast cancer cell line MCF-7 was confirmed by western blot analysis. We have also seen a correlation between bcl-2 and ER expression in human breast cancer specimens which suggests that bcl-2 could be directly involved in apoptosis associated with estrogen withdrawal. Thus we identify bcl-2 as a candidate therapeutic response variable that predicts response to tamoxifen.

Cloning and sequencing of one of the estrogen repressed clones revealed a heretofore undescribed gene that is down regulated by estrogen.

The partial sequence of the gene is as follows:

```
1  agggtgtgg gcctgtatct tgtttgctcg tcacctgtc agtcattttt ttcttcct
61  tttttaag acaaatctg aacctagaaa caccgaagcc agagcaaaaa ctatgacga
121 atccactatt tgtgcgaccc
```

Further characterization of this estrogen repressed gene is underway.

The strategy shown here could be useful in identifying genes regulated by hormones or growth factors. Clones in which genes inducible by estrogen are mutated by the U3lacZ retrovirus could be useful in screening anticancer drugs as the reporter gene lacZ faithfully follows the pattern of regulation of the endogenous genes.

### **Conclusions:**

The conclusions from project year two can be summarized as follows:

1. erbB-2 does not predict response to tamoxifen.
2. Statistical analysis of B-11 to address whether erbB-2 predicts response to chemotherapy is pending.
3. Gene trap has identified bcl-2, an inhibitor of apoptosis, as an estrogen inducible gene. Thus bcl-2 is a candidate therapeutic response variable.
4. A novel gene, which is repressed by estrogen in breast cancer cells, has been cloned using gene trap.